



TITLE:

THE LEVY LAPLACIAN AND THE LEVY PROCESS(Micro-Macro Duality in Quantum Analysis)

AUTHOR(S):

ISHIKAWA, ATSUSHI; SAITO, KIMIAKI

CITATION:

ISHIKAWA, ATSUSHI ...[et al]. THE LEVY LAPLACIAN AND THE LEVY PROCESS(Micro-Macro Duality in Quantum Analysis). 数理解析研究所講究録 2006, 1507: 14-25

ISSUE DATE:

2006-07

URL:

<http://hdl.handle.net/2433/58554>

RIGHT:

THE LÉVY LAPLACIAN AND THE LÉVY PROCESS

ATSUSHI ISHIKAWA

*Mitsubishi Electric Mechatronics Software Corporation
Nagoya 461-8670, Japan*

KIMIYAKI SAITÔ

*Department of Mathematics, Meijo University
Nagoya 468-8502, Japan
E-mail: ksaito@ccmfs.meijo-u.ac.jp*

In this paper we shall discuss the Lévy Laplacian as an operator acting on some class of the Lévy functionals. We introduce some domain of the Laplacian as a Fock space associated with the Laplacian and give associated semigroups and associated stochastic processes.

Introduction

An infinite dimensional Laplacian was introduced by P. Lévy in his famous book ¹². Since then this exotic Laplacian has been studied by many authors from various aspects see [1–6,9,10,13,15,16,17,20] and references cited therein. In this paper, we discuss a stochastic process associated with the Lévy Laplacian generalizing the methods developed in the former works [11,14,18,22–26].

This paper is organized as follows. In Section 1 we give a necessary and sufficient condition for a Fock space associated with the Lévy Laplacian. By this condition we summarize basic elements of white noise theory based on a stochastic process given as a difference of two independent Lévy processes in Section 2. In Section 3, following the recent works Kuo–Obata–Saitô ¹¹, Obata–Saitô ¹⁸, Saitô ^{23,24} and Saitô–Tsoi ²⁵, we formulate the Lévy Laplacian acting on a Hilbert space consisting of multiple Wiener integrals by the stochastic process. In Section 4, we generalize this situation by means of a direct integral of Hilbert spaces. In Section 5, based on infinitely

many Cauchy processes, we give an infinite dimensional stochastic process associated with the Lévy Laplacian.

1. Background

Let $X = \{X(t); t \in \mathbf{R}\}$ be a Lévy process, of which the characteristic function is given by

$$E[e^{izX(t)}] = \exp\{th_X(z)\}, \quad z \in \mathbf{R},$$

$$h_X(z) = imz - \frac{\sigma^2}{2}z^2 + \int_{|\lambda|>0} \left(e^{iz\lambda} - 1 - \frac{iz\lambda}{1+\lambda^2} \right) \frac{1+\lambda^2}{\lambda^2} d\beta(\lambda),$$

where $\sigma \geq 0, m \in \mathbf{R}$ and β is a positive finite measure on \mathbf{R} with $\beta(\{0\}) = \sigma^2$ and $\int_{\mathbf{R}} |\lambda|^n d\beta(\lambda) < +\infty$ for all $n \in \mathbf{N}$.

Let $E = \mathcal{S}(\mathbf{R})$ be the Schwartz space of rapidly decreasing \mathbf{R} -valued functions on \mathbf{R} . There exists an orthonormal basis $\{e_\nu\}_{\nu \geq 0}$ of $L^2(\mathbf{R})$ contained in E such that $Ae_\nu = 2(\nu + 1)e_\nu$, $\nu = 0, 1, 2, \dots$, with $A = -\frac{d^2}{du^2} + u^2 + 1$. Then by the Bochner-Minlos Theorem, there exists a probability measure μ on E^* such that

$$\int_{E^*} e^{i\langle x, \xi \rangle} d\mu(x) = \exp\left\{ \int_{\mathbf{R}} h_X(\xi(t)) dt \right\}, \quad \xi \in E.$$

Let $L^2(E^*, \mu)$ be the Hilbert space of \mathbf{C} -valued square-integrable functions on (E^*, μ) . The \mathcal{U} -transform $\mathcal{U}[\varphi]$ of $\varphi \in L^2(E^*, \mu)$ is defined by

$$\mathcal{U}[\varphi](\xi) = \exp\left\{ - \int_{\mathbf{R}} h_X(\xi(t)) dt \right\} \int_{E^*} e^{i\langle x, \xi \rangle} \varphi(x) d\mu(x), \quad \xi \in E,$$

and the Wick product $\langle \cdot, f \rangle^{\circ n}$ of $\langle \cdot, f \rangle$ is given by

$$\mathcal{U}[\langle \cdot, f \rangle^{\circ n}] = \mathcal{U}[\langle \cdot, f \rangle]^n, \quad f \in E.$$

Fixing a finite interval T on \mathbf{R} , we take an orthonormal basis $\{\zeta_n\} \subset E$ for $L^2(T)$ which is equally dense and uniformly bounded. Let $Dom(\Delta_L)$ denote the set of all $\varphi \in L^2(E^*, \mu)$ such that the limit

$$\tilde{\Delta}_L \mathcal{U}[\varphi](\xi) \equiv \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} (\mathcal{U}\varphi)''(\xi)(\zeta_n, \zeta_n),$$

exists for each $\xi \in E$ and a functional $\tilde{\Delta}_L \mathcal{U}$ is in $\mathcal{U}[(L^2)]$. The Lévy Laplacian Δ_L on $Dom(\Delta_L)$ is defined by

$$\Delta_L \varphi = \mathcal{U}^{-1} \tilde{\Delta}_L \mathcal{U} \varphi, \quad \varphi \in Dom(\Delta_L).$$

Let $F_n(\xi) = \mathcal{U}[\langle \cdot, f \rangle^{\diamond n}](\xi)$ with $\text{supp} f \subset T$. Then we can calculate that

$$F_1(\xi) = \int_{\mathbf{R}} f(t) \left(\mu + i\sigma^2 \xi(t) + \int_{|u|>0} \left(e^{i\xi(t)u} - \frac{1}{1+u^2} \right) \frac{1+u^2}{u} d\beta(u) \right) dt;$$

$$\tilde{\Delta}_L F_1(\xi) = -\frac{1}{|T|} \int_{\mathbf{R}} f(t) \int_{\mathbf{R}} u(1+u^2) e^{i\xi(t)u} d\beta(u) dt;$$

and

$$\begin{aligned} \tilde{\Delta}_L F_n(\xi) &= n F_1(\xi)^{n-1} \tilde{\Delta}_L F_1(\xi) \\ &= -\frac{n}{|T|} F_1(\xi)^{n-1} \int_{\mathbf{R}} f(t) \int_{\mathbf{R}} u(1+u^2) e^{i\xi(t)u} d\beta(u) dt. \end{aligned}$$

With these calculations, we have the following necessary and sufficient condition for a Fock space associated with the Lévy Laplacian.

Theorem 1.1. *The functionals F_n for all $n \in \mathbf{N} \cup \{0\}$ are eigenfunctions of $\tilde{\Delta}_L$ if and only if the following holds:*

- $\beta = \sigma^2 \delta_0$
- or
- $\sigma = 0, \beta = a\delta_\lambda + (a - m\lambda)\delta_{-\lambda}$ for some $\lambda > 0$ and $a \geq 0$ with $a \geq m\lambda$.

2. Preliminaries

For $p \in \mathbf{R}$ define a norm $|\cdot|_p$ by $|f|_p = |A^p f|_{L^2(\mathbf{R})}$ for $f \in E$ and let E_p be the completion of E with respect to the norm $|\cdot|_p$. Then E_p becomes a real separable Hilbert space with the norm $|\cdot|_p$ and the dual space E'_p is identified with E_{-p} by extending the inner product $\langle \cdot, \cdot \rangle$ of $L^2(\mathbf{R})$ to a bilinear form on $E_{-p} \times E_p$. It is known that $E = \text{projlim}_{p \rightarrow \infty} E_p$, and $E^* = \text{indlim}_{p \rightarrow \infty} E_{-p}$.

The canonical bilinear form on $E^* \times E$ is also denoted by $\langle \cdot, \cdot \rangle$. We denote the complexifications of $L^2(\mathbf{R})$, E and E_p by $L^2_{\mathbf{C}}(\mathbf{R})$, $E_{\mathbf{C}}$ and $E_{\mathbf{C},p}$, respectively.

Let $\{L^1_{\sigma,a,\lambda}(t)\}_{t \geq 0}$ and $\{L^2_{\sigma,a,\lambda}(t)\}_{t \geq 0}$ be independent Lévy processes of which the characteristic functions are given by

$$E[e^{izL^j_{\sigma,a,\lambda}(t)}] = e^{th(z)}, \quad t \geq 0, \quad j = 1, 2,$$

$$h(z) = imz - \frac{\sigma^2}{4}z^2 + a \left(1 + \frac{1}{\lambda^2}\right) (e^{i\lambda z} - 1),$$

where $m \in \mathbf{R}, \sigma \geq 0, a \geq 0$, and $\lambda > 0$.

Set $\Lambda_{\sigma,a,\lambda}(t) = L_{\sigma,a,\lambda}^1(t) - L_{\sigma,a,\lambda}^2(t)$ for all $t \geq 0$. Then we have

$$\begin{aligned} E[e^{iz\Lambda_{\sigma,a,\lambda}(t)}] &= e^{t(h(z)+h(-z))} \\ &= \exp \left\{ -t \frac{\sigma^2}{2} z^2 + ta \left(1 + \frac{1}{\lambda^2}\right) (e^{i\lambda z} + e^{-i\lambda z} - 2) \right\}, \quad t \geq 0. \end{aligned}$$

This characteristic function of $\Lambda_{\sigma,a,\lambda}(t)$ is corresponding to the form in Theorem 1.1 in the case of $m = 0$.

Set

$$C(\xi) = \exp \left\{ \int_{\mathbf{R}} (h(\xi_1(u)) + h(-\xi_2(u))) du \right\}, \quad \xi = (\xi_1, \xi_2) \in E \times E.$$

Then by the Bochner-Minlos Theorem, there exists a probability measure $\mu_{\sigma,a,\lambda}$ on $E^* \times E^*$ such that

$$\int_{E^* \times E^*} \exp\{i\langle x, \xi \rangle\} d\mu_{\sigma,a,\lambda}(x) = C(\xi), \quad \xi = (\xi_1, \xi_2) \in E \times E,$$

where $\langle x, \xi \rangle = \langle x_1, \xi_1 \rangle + \langle x_2, \xi_2 \rangle$, $x = (x_1, x_2) \in E^* \times E^*$, $\xi = (\xi_1, \xi_2) \in E \times E$.

Let $(L^2)_{\sigma,a,\lambda} \equiv L^2(E^* \times E^*, \mu_{\sigma,a,\lambda})$ be the Hilbert space of \mathbf{C} -valued square-integrable functions on $E^* \times E^*$ with L^2 -norm $\|\cdot\|_{\sigma,a,\lambda}$ with respect to $\mu_{\sigma,a,\lambda}$. We call an element of $(L^2)_{\sigma,a,\lambda}$ the *Lévy functional*. The Wiener-Itô decomposition theorem says that:

$$(L^2)_{\sigma,a,\lambda} = \bigoplus_{n=0}^{\infty} H_n,$$

where H_n is the space of multiple Wiener integrals of order $n \in \mathbf{N}$ and $H_0 = \mathbf{C}$. The \mathcal{U} -transform of $\varphi \in (L^2)_{\sigma,a,\lambda}$ is defined by

$$\mathcal{U}\varphi(\xi) = C(\xi)^{-1} \int_{E^* \times E^*} \varphi(x) \exp\{i\langle x, \xi \rangle\} d\mu_{\sigma,a,\lambda}(x), \quad \xi \in E \times E.$$

Theorem 2.1. ¹⁹ (see also ^{7,9,16}) *Let F be a complex-valued function defined on $E \times E$. Then F is a \mathcal{U} -transform of some Lévy functional in $(L^2)_{\sigma,a,\lambda}$ if and only if there exists a complex-valued function G defined on $E_{\mathbf{C}} \times E_{\mathbf{C}}$ such that*

- 1) for any ξ and η in $E_{\mathbf{C}} \times E_{\mathbf{C}}$, the function $G(z\xi + \eta)$ is an entire function of $z \in \mathbf{C}$,
 2) there exist nonnegative constants K and γ such that

$$|G(\xi)| \leq K \exp [\gamma |\xi|_0^2], \quad \forall \xi \in E_{\mathbf{C}} \times E_{\mathbf{C}},$$

3)

$$F(\xi) = G \left(i \frac{\sigma^2}{2} \xi_1 + i \frac{\sigma^2}{2} \xi_2 + a \left(1 + \frac{1}{\lambda^2} \right) \lambda (e^{i\lambda \xi_1} - e^{-i\lambda \xi_2}) \right)$$

for all $\xi = (\xi_1, \xi_2) \in E \times E$.

3. The Lévy Laplacian acting on the Lévy functionals

• Definition of the Lévy Laplacian

Consider $F = \mathcal{U}\varphi$ with $\varphi \in (L^2)_{\sigma,a,\lambda}$. By Theorem 2.1, for any $\xi, \eta \in E \times E$ the functions $z \mapsto F(\xi + z\eta)$ admits the Taylor series expansions:

$$F(\xi + z\eta) = \sum_{n=0}^{\infty} \frac{z^n}{n!} F^{(n)}(\xi)(\eta, \dots, \eta);$$

where $F^{(n)}(\xi) : E \times \dots \times E \rightarrow \mathbf{C}$ is a continuous n -linear functional.

Fixing a finite interval T of \mathbf{R} , we take an orthonormal basis $\{\zeta_n\}_{n=0}^{\infty} \subset E \times E$ for $L^2(T)$ which is equally dense and uniformly bounded (see e.g. 9,10). Let \mathcal{D}_L denote the set of all $\varphi \in (L^2)_{\sigma,a,\lambda}$ such that the limit

$$\tilde{\Delta}_L(\mathcal{U}\varphi)(\xi) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} (\mathcal{U}\varphi)''(\xi)(\zeta_n, \zeta_n),$$

exists for any $\xi \in E \times E$ and $\tilde{\Delta}_L(\mathcal{U}\varphi)$ is in $\mathcal{U}[(L^2)_{\sigma,a,\lambda}]$. The Lévy Laplacian Δ_L is defined by $\Delta_L \varphi = \mathcal{U}^{-1} \tilde{\Delta}_L \mathcal{U}\varphi$, $\varphi \in \mathcal{D}_L$.

• Multiple Wiener integrals by the Lévy process

Given $\sigma \geq 0, \lambda > 0, a \geq 0, n \in \mathbf{N}$ and $f \in L_{\mathbf{C}}^2(\mathbf{R})^{\otimes n}$, we consider $\varphi \in (L^2)_{\sigma,a,\lambda}$ of the form:

$$\varphi = \int_{T^n} f(u_1, \dots, u_n) d\Lambda_{\sigma,a,\lambda}(u_1) \cdots d\Lambda_{\sigma,a,\lambda}(u_n). \quad (2.1)$$

The \mathcal{U} -transform $\mathcal{U}\varphi$ of φ is given by

$$\mathcal{U}\varphi(\xi) = \int_{T^n} f(u_1, \dots, u_n) \prod_{j=1}^n \Xi_{\sigma,a,\lambda}(\xi)(u_j) du_1 \dots du_n, \quad \xi \in E \times E,$$

where

$$\Xi_{\sigma,a,\lambda}(\xi)(u_j) = i\frac{\sigma^2}{2}\xi_1(u_j) + i\frac{\sigma^2}{2}\xi_2(u_j) + a\left(1 + \frac{1}{\lambda^2}\right)\lambda(e^{i\lambda\xi_1(u_j)} - e^{-i\lambda\xi_2(u_j)}).$$

For any $\sigma \geq 0$, $\lambda > 0$, $a \geq 0$, and $n \in \mathbb{N}$ let $\mathbf{E}_{\sigma,a,\lambda,n}$ denote the space of φ which admits an expression as in (2.1), where f belongs to $L^2_{\mathbb{C}}(\mathbf{R})^{\otimes n}$ and $\text{supp } f \subset T^n$.

Set $\mathbf{E}_{\sigma,a,\lambda,0} = \mathbb{C}$ for any $\sigma \geq 0$, $\lambda > 0$, $a \geq 0$. Then $\mathbf{E}_{\sigma,a,\lambda,n}$ is a closed linear subspace of $(L^2)_{\sigma,a,\lambda}$. Using a similar method as in [25], we get the following

Theorem 3.1. *For each $\sigma \geq 0$, $n \in \mathbb{N}$, $\lambda > 0$ and $a \geq 0$ the Lévy Laplacian Δ_L becomes a scalar operator on $\mathbf{E}_{\sigma,0,\lambda,n} \cup \mathbf{E}_{0,a,\lambda,n}$ such that $\Delta_L \varphi = 0$ for all $\varphi \in \mathbf{E}_{\sigma,0,\lambda,n}$ and $\Delta_L \varphi = -\frac{n\lambda^2}{|T|} \varphi$ for all $\varphi \in \mathbf{E}_{0,a,\lambda,n}$.*

For $N \in \mathbb{N}$ and $\lambda > 0$ let $\mathbf{D}_N^{0,a,\lambda}$ be the space of $\varphi \in (L^2)_{0,a,\lambda}$ which admits an expression $\varphi = \sum_{n=1}^{\infty} \varphi_n$, $\varphi_n \in \mathbf{E}_{0,a,\lambda,n}$, such that $|||\varphi|||_{N,0,a,\lambda}^2 = \sum_{n=1}^{\infty} \alpha_N^\lambda(n) \|\varphi_n\|_{0,a,\lambda}^2 < \infty$, where $\alpha_N^\lambda(n) = \sum_{\ell=0}^N \left(\frac{n\lambda^2}{|T|}\right)^{2\ell}$.

By the Schwartz inequality we see that $\mathbf{D}_N^{0,a,\lambda}$ is a subspace of $(L^2)_{0,a,\lambda}$ and becomes a Hilbert space equipped with the new norm $||| \cdot |||_{N,0,a,\lambda}$.

Moreover, in view of the inclusion relations:

$$(L^2)_{0,a,\lambda} \supset \mathbf{D}_1^{0,a,\lambda} \supset \dots \supset \mathbf{D}_N^{0,a,\lambda} \supset \mathbf{D}_{N+1}^{0,a,\lambda} \supset \dots,$$

we define

$$\mathbf{D}_\infty^{0,a,\lambda} = \text{proj} \lim_{N \rightarrow \infty} \mathbf{D}_N^{0,a,\lambda} = \bigcap_{N=1}^{\infty} \mathbf{D}_N^{0,a,\lambda}.$$

Then Δ_L becomes a continuous linear operator defined on $\mathbf{D}_{N+1}^{0,a,\lambda}$ into $\mathbf{D}_N^{0,a,\lambda}$ satisfying $|||\Delta_L \varphi|||_{N,0,a,\lambda} \leq |||\varphi|||_{N+1,0,a,\lambda}$, $\varphi \in \mathbf{D}_\infty^{0,a,\lambda}$, $N \in \mathbb{N}$. Therefore Δ_L is a continuous linear operator on $\mathbf{D}_\infty^{0,a,\lambda}$. Moreover the operator Δ_L is a self-adjoint operator densely defined in $\mathbf{D}_N^{0,a,\lambda}$ for each $N \in \mathbb{N}$ and $\lambda > 0$.

For each $t \geq 0$, $\lambda > 0$ and $a \geq 0$ we consider an operator G_t^λ on $\mathbf{D}_\infty^{0,a,\lambda}$ defined by

$$G_t^\lambda \varphi = \sum_{n=1}^{\infty} e^{-tn\lambda^2/|T|} \varphi_n, \quad \varphi = \sum_{n=1}^{\infty} \varphi_n \in \mathbf{D}_\infty^{0,a,\lambda}.$$

We also define G_t^0 on $(L^2)_{\sigma,0,\lambda}$ as an identity operator by

$$G_t^0 \varphi = \varphi, \quad \varphi \in (L^2)_{\sigma,0,\lambda}.$$

Theorem 3.2. ^{11,23} *Let $\lambda > 0$ and $a \geq 0$. Then the family of operators $\{G_t^\lambda; t \geq 0\}$ on $\mathbf{D}_\infty^{0,a,\lambda}$ is an equi-continuous semigroup of class (C_0) of which the infinitesimal generator is Δ_L .*

4. Extensions of the Lévy Laplacian

Let $d\nu(\lambda)$ be a finite Borel measure on \mathbf{R} satisfying

$$\int_{(0,\infty)} \frac{d\nu(\lambda)}{\lambda^4} < \infty.$$

Fix $N \in \mathbf{N}$ and $a \geq 0$. Let \mathfrak{D}_N^σ be the space of (equivalent classes of) measurable vector functions $\varphi = (\varphi^\lambda)$ with $\varphi^\lambda = \sum_{n=1}^\infty \varphi_n^\lambda \in \mathbf{D}_N^{0,a,\lambda}$ for all $\lambda > 0$, and $\varphi^0 \in (L^2)_{\sigma,0,\lambda}$, such that

$$|||\varphi|||_N^2 = \|\varphi^0\|_{\sigma,0,\lambda}^2 + \sum_{n=1}^\infty \int_{(0,\infty)} \|\varphi_n^\lambda\|_{0,a,\lambda,N}^2 d\nu(\lambda) < \infty. \quad (3.1)$$

Then \mathfrak{D}_N^σ becomes a Hilbert space with the norm given in (3.1).

In view of the natural inclusion $\mathfrak{D}_{N+1}^\sigma \subset \mathfrak{D}_N^\sigma$ for $N \in \mathbf{N}$, which is obvious from construction, we define $\mathfrak{D}_\infty^\sigma = \text{proj} \lim_{N \rightarrow \infty} \mathfrak{D}_N^\sigma = \bigcap_{N=1}^\infty \mathfrak{D}_N^\sigma$.

The Lévy Laplacian Δ_L is defined on the space $\mathfrak{D}_\infty^\sigma$ by

$$\Delta_L \varphi = (\Delta_L \varphi^\lambda), \quad \varphi = (\varphi^\lambda) \in \mathfrak{D}_\infty^\sigma.$$

Then Δ_L is a continuous linear operator from $\mathfrak{D}_\infty^\sigma$ into itself. Similarly, for $t \geq 0$ we define

$$G_t \varphi = (G_t^\lambda \varphi^\lambda), \quad \varphi = (\varphi^\lambda) \in \mathfrak{D}_\infty^\sigma.$$

Then by Theorem 3.2 we have the following:

Theorem 4.1. *The family $\{G_t; t \geq 0\}$ is an equi-continuous semigroup of class (C_0) on $\mathfrak{D}_\infty^\sigma$ whose generator is given by Δ_L .*

Remark: Let \tilde{G}_t be an operator defined on $\mathcal{U}[\mathfrak{D}_\infty^\sigma]$ by $\tilde{G}_t = \mathcal{U} G_t \mathcal{U}^{-1}$, $t \geq 0$. Then by the above theorem, $\{\tilde{G}_t; t \geq 0\}$ is an equi-continuous semigroup of class (C_0) generated by the operator $\tilde{\Delta}_L$.

5. Associated infinite dimensional stochastic processes

• Space $E^{[0,\infty)}$

For $p \in \mathbf{R}$ let $E_p^{[0,\infty)}$ be the linear space of all functions $\lambda \mapsto \xi_\lambda \in E_p \times E_p$, $\lambda \geq 0$, which are strongly measurable. An element of $E_p^{[0,\infty)}$ is denoted by $\xi = (\xi_\lambda)_{\lambda \geq 0}$. Equipped with the metric given by

$$d_p(\xi, \eta) = \int_{[0,\infty)} \frac{|\xi_\lambda - \eta_\lambda|_p}{1 + |\xi_\lambda - \eta_\lambda|_p} d\nu(\lambda), \quad \xi = (\xi_\lambda), \quad \eta = (\eta_\lambda),$$

the space $E_p^{[0,\infty)}$ becomes a complete metric space.

In view of $d_p \leq d_q$ for $p \geq q$, we introduce the projective limit space

$$E^{[0,\infty)} = \text{proj} \lim_{p \rightarrow \infty} E_p^{[0,\infty)}.$$

• Space $\mathbf{C}^{[0,\infty)}$

Similarly, let $\mathbf{C}^{[0,\infty)}$ denote the linear space of all measurable function $\lambda \mapsto z_\lambda \in \mathbf{C}$ equipped with the metric defined by

$$\rho(\mathbf{z}, \mathbf{u}) = \int_{[0,\infty)} \frac{|z_\lambda - u_\lambda|}{1 + |z_\lambda - u_\lambda|} d\nu(\lambda), \quad \mathbf{z} = (z_\lambda), \quad \mathbf{u} = (u_\lambda).$$

Then $\mathbf{C}^{[0,\infty)}$ is also a complete metric space.

• Extension of the \mathcal{U} -transform

The \mathcal{U} -transform can be extended to a continuous linear operator on $\mathcal{D}_\infty^\sigma$ by

$$\mathcal{U}\varphi(\xi) = (\mathcal{U}\varphi^\lambda(\xi_\lambda))_{\lambda \geq 0}, \quad \xi = (\xi_\lambda)_{\lambda \geq 0} \in E^{[0,\infty)},$$

for any $\varphi = (\varphi^\lambda)_{\lambda \geq 0} \in \mathcal{D}_\infty^\sigma$.

The space $\mathcal{U}[\mathcal{D}_\infty^\sigma]$ is endowed with the topology induced from $\mathcal{D}_\infty^\sigma$ by the \mathcal{U} -transform. Then the \mathcal{U} -transform becomes a homeomorphism from $\mathcal{D}_\infty^\sigma$ onto $\mathcal{U}[\mathcal{D}_\infty^\sigma]$. The transform $\mathcal{U}\varphi$ of $\varphi \in \mathcal{D}_\infty^\sigma$ is a continuous operator from $E^{[0,\infty)}$ into $\mathbf{C}^{[0,\infty)}$. We denote the operator by the same notation $\mathcal{U}\varphi$.

• Associated stochastic process

Let $\{X_t^j\}, j = 1, 2$, be independent Cauchy processes with t running over $[0, \infty)$, of which the characteristic functions are given by

$$\mathbf{E}[e^{izX_t^j}] = e^{-t|z|}, \quad z \in \mathbf{R}, \quad j = 1, 2.$$

Take a smooth function $\eta_T \in E$ with $\eta_T = 1/|T|$ on T .

Set

$$Y_t^\lambda = (X_{\lambda t}^1 \eta_T, -X_{\lambda t}^2 \eta_T) \quad \lambda \geq 0.$$

Define an infinite dimensional stochastic process $\{\mathbf{Y}_t; t \geq 0\}$ starting at $\xi = (\xi_\lambda)_{\lambda \geq 0} \in E^{[0, \infty)}$ by

$$\mathbf{Y}_t = (\xi_\lambda + Y_t^\lambda)_{\lambda \geq 0}, \quad t \geq 0.$$

Then this is an $E^{[0, \infty)}$ -valued stochastic process and we have the following

Theorem 5.1. *If F is the \mathcal{U} -transform of an element in $\mathfrak{D}_\infty^\sigma$, we have*

$$\tilde{G}_t F(\xi) = E[F(\mathbf{Y}_t) | \mathbf{Y}_0 = \xi], \quad t \geq 0. \quad (4.1)$$

Proof. We first consider the case when $F \in \mathcal{U}[\mathfrak{D}_\infty^\sigma]$ is given by

$$F(\xi) = (F^\lambda(\xi_\lambda))_{\lambda \geq 0},$$

$$F^0 \in \mathcal{U}[(L^2)_{\sigma, 0, \lambda}],$$

$$F^\lambda(\xi_\lambda) = \left(a\lambda \left(1 + \frac{1}{\lambda^2} \right) \right)^n \int_{T^n} f(\mathbf{u}) \cdot \prod_{j=1}^n \left\{ e^{i\lambda \xi_{1, \lambda}(u_j)} - e^{-i\lambda \xi_{2, \lambda}(u_j)} \right\} d\mathbf{u},$$

with $f \in L^2_{\mathbf{C}}(\mathbf{R})^{\otimes n}$. Then we have

$$\begin{aligned} E[F(\mathbf{Y}_t) | \mathbf{Y}_0 = \xi] &= (E[F^\lambda(\xi_\lambda + Y_t^\lambda)])_{\lambda \geq 0} \\ &= \left(F^0(\xi_0) \delta_{\lambda, 0} + \left(a\lambda \left(1 + \frac{1}{\lambda^2} \right) \right)^n \int_{T^n} f(\mathbf{u}) E \left[\prod_{j=1}^n \left\{ e^{i\lambda \xi_{1, \lambda}(u_j)} e^{i \frac{\lambda}{|T|} X_{\lambda t}^1 1_{(0, \infty)}(\lambda)} \right. \right. \right. \\ &\quad \left. \left. \left. - e^{-i\lambda \xi_{2, \lambda}(u_j)} e^{i \frac{\lambda}{|T|} X_{\lambda t}^2 1_{(0, \infty)}(\lambda)} \right\} \right] d\mathbf{u} \right)_{\lambda \geq 0} \\ &= \left(e^{-tn\lambda/|T|} F^\lambda(\xi_\lambda) \right)_{\lambda \geq 0} = (\tilde{G}_t^\lambda F^\lambda(\xi_\lambda))_{\lambda \geq 0} = \tilde{G}_t F(\xi). \end{aligned}$$

Next let $F = (F^0 \delta_{\lambda,0} + \sum_{n=1}^{\infty} F_n^\lambda)_{\lambda \geq 0} \in \mathcal{U}[\mathcal{D}_\infty^\sigma]$. Then for ν -almost all $\lambda > 0$ and for any $n \in \mathbb{N}$, F_n^λ is expressed in the following form:

$$F_n^\lambda(\xi_\lambda) = \lim_{N \rightarrow \infty} \left(a\lambda \left(1 + \frac{1}{\lambda^2} \right) \right)^n \int_{T^n} f_\lambda^{[N]}(\mathbf{u}) \prod_{j=1}^n \left\{ e^{i\lambda \xi_{1,\lambda}(u_j)} - e^{-i\lambda \xi_{2,\lambda}(u_j)} \right\} d\mathbf{u}.$$

Since $F^0 \in \mathcal{U}[(L^2)_{\sigma,0,\lambda}]$ and $F_n^\lambda \in \mathcal{U}[\mathbf{D}_\infty^{0,a,\lambda}]$, there exist $\varphi^0 \in (L^2)_{\sigma,0,\lambda}$ and $\varphi_n^\lambda \in \mathbf{D}_\infty^{0,a,\lambda}$ such that $F^0 = \mathcal{U}[\varphi^0]$ and $F_n^\lambda = \mathcal{U}[\varphi_n^\lambda]$ for ν -almost all λ and each n . By the Schwarz inequality, we see that

$$\begin{aligned} \sum_{n=1}^{\infty} E[|F_n^\lambda(\xi_\lambda + Y_t^\lambda)|] &\leq \sum_{n=0}^{\infty} \|\varphi_n^\lambda\|_{0,a,\lambda} E[\|\varphi_{\xi_\lambda + Y_t^\lambda}\|_{0,a,\lambda}] \\ &\leq \left\{ \sum_{n=1}^{\infty} \alpha_N^\lambda(n)^{-1} \right\}^{1/2} \left\{ \sum_{n=1}^{\infty} \alpha_N^\lambda(n) \|\varphi_n^\lambda\|_{0,a,\lambda}^2 \right\}^{1/2} E[\|\varphi_{\xi_\lambda + Y_t^\lambda}\|_{0,a,\lambda}] < \infty, \end{aligned}$$

where $\varphi_{\xi_\lambda} = C(\xi_\lambda)^{-1} e^{i\langle \cdot, \xi_\lambda \rangle}$ for ν -almost all $\lambda \geq 0$ and each $N \in \mathbb{N}$. Therefore by the continuity of $\widetilde{G}_t^\lambda, \lambda \geq 0$, we get that

$$\begin{aligned} E[F(\xi + \mathbf{Y}_t)] &= \left(\sum_{n=1}^{\infty} E[F_n^\lambda(\xi_\lambda + Y_t^\lambda)] \right)_{\lambda \geq 0} = \left(\sum_{n=1}^{\infty} \widetilde{G}_t^\lambda F_n^\lambda(\xi_\lambda) \right)_{\lambda \geq 0} \\ &= \left(\widetilde{G}_t^\lambda \sum_{n=1}^{\infty} F_n^\lambda(\xi_\lambda) \right)_{\lambda \geq 0} = \widetilde{G}_t F(\xi). \end{aligned}$$

Thus we obtain the assertion. \square

For any $\eta \in E^{[0,\infty)}$ we define a translation operator T_η on $\mathcal{D}_\infty^\sigma$ by

$$(\mathcal{U}T_\eta \varphi)(\xi) = (\mathcal{U}\varphi)(\xi + \eta).$$

Theorem 5.2. *For all φ in $\mathcal{D}_\infty^\sigma$ we have*

$$G_t \varphi = E[T_{(Y_t^\lambda)_{\lambda > 0}} \varphi], \quad t \geq 0. \quad (4.2)$$

Acknowledgements

The authors would like to express their deep thanks to Professor I. Ojima, RIMS, Kyoto University, for his hard work of organizing the stimulating workshop.

References

1. L. Accardi and V. Bogachëv: *The Ornstein-Uhlenbeck process associated with the Lévy Laplacian and its Dirichlet form*, Prob. Math. Stat. **17** (1997), 95–114.
2. L. Accardi, P. Gibilisco and I. V. Volovich: *Yang-Mills gauge fields as harmonic functions for the Lévy Laplacian*, Russ. J. Math. Phys. **2** (1994), 235–250.
3. L. Accardi and O. G. Smolyanov: *Trace formulae for Levy-Gaussian measures and their application*, in “Mathematical Approach to Fluctuations Vol. II (T. Hida, Ed.),” pp. 31–47, World Scientific, 1995.
4. D. M. Chung, U. C. Ji and K. Saitô: *Cauchy problems associated with the Lévy Laplacian in white noise analysis*, Infin. Dimen. Anal. Quantum Probab. Rel. Top. **2** (1999), 131–153.
5. M. N. Feller: *Infinite-dimensional elliptic equations and operators of Lévy type*, Russ. Math. Surveys **41** (1986), 119–170.
6. T. Hida: “Analysis of Brownian Functionals,” Carleton Math. Lect. Notes, No. 13, Carleton University, Ottawa, 1975.
7. T. Hida, H.-H. Kuo, J. Potthoff and L. Streit: “White Noise: An Infinite Dimensional Calculus,” Kluwer Academic, 1993.
8. I. Kubo and S. Takenaka: *Calculus on Gaussian white noise I–IV*, Proc. Japan Acad. **56A** (1980) 376–380; **56A** (1980) 411–416; **57A** (1981) 433–436; **58A** (1982) 186–189.
9. H.-H. Kuo: “White Noise Distribution Theory,” CRC Press, 1996.
10. H.-H. Kuo, N. Obata and K. Saitô: *Lévy Laplacian of generalized functions on a nuclear space*, J. Funct. Anal. **94** (1990), 74–92.
11. H.-H. Kuo, N. Obata and K. Saitô: *Diagonalization of the Lévy Laplacian and related stable processes*, Infin. Dimen. Anal. Quantum Probab. Rel. Top. **5** (2002), 317–331.
12. P. Lévy: “Leçons d’Analyse Fonctionnelle,” Gauthier-Villars, Paris, 1922.
13. R. Léandre and I. A. Volovich: *The stochastic Lévy Laplacian and Yang-Mills equation on manifolds*, Infin. Dimen. Anal. Quantum Probab. Rel. Top. **4** (2001) 161–172.
14. K. Nishi and K. Saitô: *An infinite dimensional stochastic process and the Lévy Laplacian acting on WND-valued functions*, “Quantum Information and Complexity” pp. 376–390, World Scientific, 2004.
15. N. Obata: *A characterization of the Lévy Laplacian in terms of infinite dimensional rotation groups*, Nagoya Math. J. **118** (1990), 111–132.
16. N. Obata: “White Noise Calculus and Fock Space,” Lect. Notes in Math. Vol. 1577, Springer-Verlag, 1994.
17. N. Obata: *Quadratic quantum white noises and Lévy Laplacian*, Nonlinear Analysis **47** (2001), 2437–2448.
18. N. Obata and K. Saitô: *Cauchy processes and the Lévy Laplacian*, Quantum Probability and White Noise Analysis **16** (2002), 360–373.
19. J. Potthoff and L. Streit: *A characterization of Hida distributions*, J. Funct. Anal. **101** (1991), 212–229.

20. K. Saitô: *Itô's formula and Lévy's Laplacian I*, Nagoya Math. J. **108** (1987), 67–76; *II*, *ibid.* **123** (1991), 153–169.
21. K. Saitô: *A (C_0) -group generated by the Lévy Laplacian II*, *Infin. Dimen. Anal. Quantum Probab. Rel. Top.* **1** (1998) 425–437.
22. K. Saitô: *A stochastic process generated by the Lévy Laplacian*, *Acta Appl. Math.* **63** (2000), 363–373.
23. K. Saitô: *The Lévy Laplacian and stable processes*, *Chaos, Solitons and Fractals* **12** (2001), 2865–2872.
24. K. Saitô: *An infinite dimensional Laplacian acting on multiple Wiener integrals by some Lévy processes*, to appear in “Infinite Dimensional Harmonic Analysis” **III** (2005).
25. K. Saitô and A. H. Tsoi: *The Lévy Laplacian as a self-adjoint operator*, in “Quantum Information,” pp. 159–171, World Scientific, 1999.
26. K. Saitô and A. H. Tsoi: *The Lévy Laplacian acting on Poisson noise functionals*, *Infin. Dimen. Anal. Quantum Probab. Rel. Top.* **2** (1999), 503–510.